

REPORT
OF THE
CANADIAN ARCTIC EXPEDITION
1913-18

VOLUME VII: CRUSTACEA

PART E: AMPHIPODS

By CLARENCE R. SHOFMAKER



OTTAWA
THOMAS MULVEY
PRINTER TO THE KING'S MOST EXCELLENT MAJESTY
1920

Vol. vii—67920—1

Issued Sept. 7th, 1920

1991 1992 1993 1994 1995 1996 1997 1998 1999 2000 2001 2002 2003 2004 2005 2006 2007 2008 2009 2010 2011 2012 2013 2014 2015 2016 2017 2018 2019 2020 2021 2022 2023 2024 2025 2026 2027 2028 2029 2030 2031 2032 2033 2034 2035 2036 2037 2038 2039 2040 2041 2042 2043 2044 2045 2046 2047 2048 2049 2050 2051 2052 2053 2054 2055 2056 2057 2058 2059 2060 2061 2062 2063 2064 2065 2066 2067 2068 2069 2070 2071 2072 2073 2074 2075 2076 2077 2078 2079 2080 2081 2082 2083 2084 2085 2086 2087 2088 2089 2090 2091 2092 2093 2094 2095 2096 2097 2098 2099 2100 2101 2102 2103 2104 2105 2106 2107 2108 2109 2110 2111 2112 2113 2114 2115 2116 2117 2118 2119 2120 2121 2122 2123 2124 2125 2126 2127 2128 2129 2130 2131 2132 2133 2134 2135 2136 2137 2138 2139 2140 2141 2142 2143 2144 2145 2146 2147 2148 2149 2150 2151 2152 2153 2154 2155 2156 2157 2158 2159 2160 2161 2162 2163 2164 2165 2166 2167 2168 2169 2170 2171 2172 2173 2174 2175 2176 2177 2178 2179 2180 2181 2182 2183 2184 2185 2186 2187 2188 2189 2190 2191 2192 2193 2194 2195 2196 2197 2198 2199 2200 2201 2202 2203 2204 2205 2206 2207 2208 2209 2210 2211 2212 2213 2214 2215 2216 2217 2218 2219 2220 2221 2222 2223 2224 2225 2226 2227 2228 2229 2230 2231 2232 2233 2234 2235 2236 2237 2238 2239 2240 2241 2242 2243 2244 2245 2246 2247 2248 2249 2250 2251 2252 2253 2254 2255 2256 2257 2258 2259 2260 2261 2262 2263 2264 2265 2266 2267 2268 2269 2270 2271 2272 2273 2274 2275 2276 2277 2278 2279 2280 2281 2282 2283 2284 2285 2286 2287 2288 2289 2290 2291 2292 2293 2294 2295 2296 2297 2298 2299 2300 2301 2302 2303 2304 2305 2306 2307 2308 2309 2310 2311 2312 2313 2314 2315 2316 2317 2318 2319 2320 2321 2322 2323 2324 2325 2326 2327 2328 2329 2330 2331 2332 2333 2334 2335 2336 2337 2338 2339 2340 2341 2342 2343 2344 2345 2346 2347 2348 2349 2350 2351 2352 2353 2354 2355 2356 2357 2358 2359 2360 2361 2362 2363 2364 2365 2366 2367 2368 2369 2370 2371 2372 2373 2374 2375 2376 2377 2378 2379 2380 2381 2382 2383 2384 2385 2386 2387 2388 2389 2390 2391 2392 2393 2394 2395 2396 2397 2398 2399 2400 2401 2402 2403 2404 2405 2406 2407 2408 2409 2410 2411 2412 2413 2414 2415 2416 2417 2418 2419 2420 2421 2422 2423 2424 2425 2426 2427 2428 2429 2430 2431 2432 2433 2434 2435 2436 2437 2438 2439 2440 2441 2442 2443 2444 2445 2446 2447 2448 2449 2450 2451 2452 2453 2454 2455 2456 2457 2458 2459 2460 2461 2462 2463 2464 2465 2466 2467 2468 2469 2470 2471 2472 2473 2474 2475 2476 2477 2478 2479 2480 2481 2482 2483 2484 2485 2486 2487 2488 2489 2490 2491 2492 2493 2494 2495 2496 2497 2498 2499 2500 2501 2502 2503 2504 2505 2506 2507 2508 2509 2510 2511 2512 2513 2514 2515 2516 2517 2518 2519 2520 2521 2522 2523 2524 2525 2526 2527 2528 2529 2530 2531 2532 2533 2534 2535 2536 2537 2538 2539 2540 2541 2542 2543 2544 2545 2546 2547 2548 2549 2550 2551 2552 2553 2554 2555 2556 2557 2558 2559 2560 2561 2562 2563 2564 2565 2566 2567 2568 2569 2570 2571 2572 2573 2574 2575 2576 2577 2578 2579 2580 2581 2582 2583 2584 2585 2586 2587 2588 2589 2590 2591 2592 2593 2594 2595 2596 2597 2598 2599 2600 2601 2602 2603 2604 2605 2606 2607 2608 2609 2610 2611 2612 2613 2614 2615 2616 2617 2618 2619 2620 2621 2622 2623 2624 2625 2626 2627 2628 2629 2630 2631 2632 2633 2634 2635 2636 2637 2638 2639 2640 2641 2642 2643 2644 2645 2646 2647 2648 2649 2650 2651 2652 2653 2654 2655 2656 2657 2658 2659 2660 2661 2662 2663 2664 2665 2666 2667 2668 2669 2670 2671 2672 2673 2674 2675 2676 2677 2678 2679 2680 2681 2682 2683 2684 2685 2686 2687 2688 2689 2690 2691 2692 2693 2694 2695 2696 2697 2698 2699 2700 2701 2702 2703 2704 2705 2706 2707 2708 2709 2710 2711 2712 2713 2714 2715 2716 2717 2718 2719 2720 2721 2722 2723 2724 2725 2726 2727 2728 2729 2730 2731 2732 2733 2734 2735 2736 2737 2738 2739 2740 2741 2742 2743 2744 2745 2746 2747 2748 2749 2750 2751 2752 2753 2754 2755 2756 2757 2758 2759 2760 2761 2762 2763 2764 2765 2766 2767 2768 2769 2770 2771 2772 2773 2774 2775 2776 2777 2778 2779 2780 2781 2782 2783 2784 2785 2786 2787 2788 2789 2790 2791 2792 2793 2794 2795 2796 2797 2798 2799 2800 2801 2802 2803 2804 2805 2806 2807 2808 2809

The Amphipods of the Canadian Arctic Expedition, 1913-18.

By CLARENCE R. SHOEMAKER

Division of Marine Invertebrates, U. S. National Museum

The marine and freshwater amphipods collected by the naturalist of the southern party of the Canadian Arctic Expedition, Mr. Frits Johansen, are for the most part well-known arctic forms, but one, a species of *Synurella*, is new to science, this genus now appearing for the first time in American waters.

Kutius obesus was known from the Atlantic only on the strength of two specimens, until thirty-five were taken off the southwestern coast of Greenland by the *Tjalfe* Expedition in 1908-9; it appears now for the first time in the Pacific.

The known ranges of several species have been greatly extended by the records in the collection under consideration; the details will be given under the species involved.

The collection contains fifty-three species included in forty-one genera which are distributed among eighteen families; the family Lysianassidae, as would be expected in an arctic collection, is represented by the greatest number of genera, species, and individuals.

An appendix has been added consisting of data based upon specimens from the *Neptune* and other Canadian expeditions.

The color notes given under some of the species are based upon colored sketches made by Mr. Frits Johansen. The color nomenclature is based upon Ridgway's "Color Standards and Nomenclature."

Order AMPHIPODA.

SUB-ORDER GAMMARIDEA.

Family LYSIANASSIDÆ.

1. *Anonyx nugax* (Phipps).

1774. *Cancer nugax* PHIPPS, Voy. North Pole, p. 192, pl. 12, f. 2.
1906. *Anonyx nugax* + *A. lagena* STEBBING, Tierreich, Amph. I, p. 54, and synonymy.
1911. *Anonyx nugax* STAPPERS, Crust. Malacost., Campagne Arctique de 1907, du Due d'Orléans, p. 8.
1913. *Anonyx nugax* STEPHENSEN, Conspectus Crust. et Pycnog. Groenl., p. 115.

Station 29f: 70° 13' N., 140° 50' W., April 4, 1914, from stomach of *Phoca hispida* Schreber, water depth about 30 fathoms; 6 specimens.

Station 43a: Dolphin and Union strait (off Cockburn point), Northwest Territories, September 13, 1915, 50 fathoms, mud with pebbles, but no algæ; 2 specimens.

Station 46e: Dolphin and Union strait (off Bernard harbour), Northwest Territories, February 16, 1916, 6 fathoms; 300 specimens.

Station 46g: Dolphin and Union strait (Bernard harbour), Northwest Territories, May 4, 1916; from Eskimos; 7 specimens.

Distribution.—This species is very widely distributed, being found throughout the Arctic, North Atlantic, and North Pacific oceans.

2. *Hippomedon holbølli* (Krøyer).

1846. *Anonyx holbølli* KRØYER, Naturh. Tidsskr., ser. 2, vol. 2, p. 8, 38.
1906. *Hippomedon holbølli* STEBBING, Tierreich, Amph. I, p. 58, and synonymy.
1911. *Hippomedon holbølli* STAPPERS, Crust. Malacost., Campagne Arctique de 1907, du Due d'Orléans, p. 6.

Station 29f: 70° 13' N., 140° 50' W., April 4, 1914. From stomach of *Phoca hispida* Schreber; water depth about 30 fathoms; 30 specimens.
Distribution.—Arctic ocean.

3. *Onisimus botkini* Birula.

1897. *Onisimus botkini* BIRULA, Annuaire du Musée Zoologique de l'Acad. Imp. des Sciences de St. Pétersbourg, vol. II, 1897, p. 105.

1900. *Onisimus botkini* BRÜGGEN, Mém. de l'Acad. Imp. des Sciences de St. Pétersbourg, sér. 8, vol. 18, No. 16, p. 7, pl. II, f. 20-25.

Station 27c: Lagoon-bay at Collinson point, Alaska, September 15, 1913, 0-1 foot water; 1 specimen.

Station 28o: Collinson point, Alaska, July 8, 1914, from stomach of *Salvelinus malma* Walb.; 7 specimens.

Station 28r: Bay at Collinson Point, Alaska, July 24, 1914; from stomach of *Cottus quadricornis* L., 3 fragments.

Station 30c: Demarcation point, Alaska, May 10, 1914, 3 fathoms, sandy mud; 200 specimens.




Figure 1. *Onisimus botkini* Birula ♂.

1. Head and antennae. 2 Maxilla 1. 3 Mandible. 4 2d and 3d abdominal segments and urosome. 5 Telson, and 2d and 3d uropods.

The specimens examined by Birula and by von der Brüggen all came from shallow water in the Kara sea, so it is very interesting to see these very fine specimens from Alaska. As these specimens differ in a few minor details from those collected in the Kara sea, I here add a brief description and a few figures.




Figure 2. *Onisimus bolkini* Birula ♂.

1 Peraeopod 2. 2 Gnathopod 2. 3 Gnathopod 1. 4 Peraeopod 5

Head; lateral lobes narrowly rounded. Eyes; pear-shaped, red. Antenna 1; 1st joint about twice as long as 2nd and 3rd combined, slightly produced at the upper, inside, anterior edge, and provided on the upper, inside, posterior end with a few plumose setae, flagellum in ♂, 24 to 30 joints, in ♀, 14 to 18 joints, accessory flagellum, 5-7 joints, of which the first joint is about as long as all the rest combined. Antenna 2 not much longer than 1, ultimate joint slightly shorter than penultimate, flagellum in ♂, 30 to 38 joints, in ♀, 18 to 23 joints.

Side-plate 1 expanded below, the lower anterior corner produced and broadly rounded, the lower edge nearly straight, a slight notch bearing a minute seta at lower posterior corner. Side-plates 2 and 3 with the sides nearly parallel, slightly wider below, the lower edge nearly straight. Side-plate 4 deeply and broadly emarginate behind, the lower part wide and the lower edge nearly straight.

Gnathopod 1, 6th joint about one-third longer than 5th with front edge slightly convex and back edge slightly concave, palm oblique and provided throughout its entire length with a single row of fine, sharp teeth and several spines, dactyl curved, with a single small tooth or spine in the centre of the inside edge, back of this tooth a row of very fine serrations, 4th and 5th joints provided on their under surface with mats of fine setae. Gnathopod 2 as in *O. edwardsii* or *O. plautus*, except that 5th and 6th joints are much more setose.

The pereopods are rather short and stout. Third abdominal segment produced at the lower posterior corner into a small rounded lobe which varies much in size.

Telson a little longer than wide, sides convex, posterior border very slightly emarginate and bearing two minute setae.

The largest specimens measure about 19 mm.

4. *Onisimus brevicaudatus* Hansen.

1886. *Onisimus brevicaudatus* HANSEN, *Dijmphna-Togteta* Ubd., p. 216, pl. 21, f. 7-7e.
1906. *Onisimus brevicaudatus* STEBBING, *Tierreich*, Amph. I, p. 27, and synonymy.

Station 29f: Lat. 70° 13' N., long. 140° 50' W., April 4, 1914, from stomach of *Phoca hispida* Schreber. Water depth about 30 fathoms. 6 specimens.
The specimens which Hansen described came from the Kara sea.

5. *Onisimus plautus* (Krøyer).

1845. *Anonyx plautus* KRØYER, *Naturh. Tidsskr.*, ser. 2, vol. I, p. 629.
1906. *Onisimus plautus* STEBBING, *Tierreich*, Amph. I, p. 26, and synonymy.
1911. *Onisimus plautus* STAPPERS, *Crust. Malacost.*, Campagne Arctique de 1907, du Oue d'Orleans, p. 16.

Station 46c: Dolphin and Union strait, Northwest Territories (off Bernard harbour), pelagic, over 38 feet water; February 19, 1916; 3 specimens.

Distribution.—Arctic ocean, North Atlantic, and North sea, Skagerrak (Bohuslän).

6. *Orchomenella groenlandica* (Hansen).

1887. *Anonyx groenlandicus* HANSEN, *Vid. Meddel.*, ser. 4, vol. 9, p. 72, pl. 2, f. 5-5g.
1906. *Orchomenella groenlandica* STEBBING, *Tierreich*, Amph. I, p. 83, and synonymy.
1913. *Orchomenella groenlandica* STEPHENSEN, *Meddel. om Grønland*, vol. XXII, p. 123.

Station 41: Bernard harbour, Northwest Territories (outer harbour), July 20, 1915, 5 fathoms, sandy mud with algæ; 1 specimen.

Distribution.—Arctic ocean, East Greenland, Finland.

7. *Orchomenella minuta* (Krøyer).

1846. *Anonyx minutus* KRØYER, Naturh. Tidsskr., ser. 2, vol. 2, p. 23.
 1906. *Orchomenella minuta* STEBBING, Tierreich, Amph. I, p. 82, and synonymy.
 1909. *Orchomenella minuta* BRÜGGEN, Mém. Acad. Imp. Sci. St. Pétersbourg, sér. 5, vol. 18, No. 16, p. 13.
 1916. *Orchomenella minuta* STEPHENSEN, Meddel. om Grønland, vol. LIII, p. 285.

Station 37j: Bernard harbour, Northwest Territories, September 1, 1914, pelagic, over 2 fathoms of water; 2 specimens.

Station 46c: Dolphin and Union strait, Northwest Territories, February 19, 1916, pelagic, over 38 feet water (off Bernard harbour), 1 specimen.

Distribution.—Arctic Ocean, North Atlantic, North Sea, Greenland.

8. *Pseudalibrotus glacialis* G. O. Sars.

1900. *Pseudalibrotus glacialis* G. O. Sars, in NANSEN, Norwegian North Polar Exp., vol. I, No. 5, p. 31, pl. 6.

Station 28o: Collinson point, Alaska, July 8, 1914, from stomach of *Salvelinus malma* Walb.; 7 specimens.

Station 57a: Cape Smyth (point Barrow), Alaska, August 8, 1916, pelagic, over 1 fathom of water; 1 specimen.

The specimens described by Sars were obtained north of Franz Josef land in 1894. The present specimens, coming from Alaska, indicate a wide distribution for the species.

9. *Pseudalibrotus litoralis* (Krøyer).

1845. *Anonyx litoralis* KRØYER, Naturh. Tidsskr., ser. 2, vol. I, p. 621.
 1906. *Pseudalibrotus litoralis* STEBBING, Tierreich, Amph. I, p. 33.
 1911. *Pseudalibrotus litoralis* STAPPERS, Crust. Malacost., Campagne Arctique de 1907 du Duc d'Orleans, p. 14.

Station 27h: Lagoon-bay at Collinson point, Alaska, September 18, 1913, 0-1 foot water; 1 specimen.

Station 28o: Collinson point, Alaska, July 8, 1914, from stomach of *Salvelinus malma* Walb.; 10 specimens.

Stations 37r, u: Bernard harbour, Northwest Territories, October 16-20, 1915, pelagic, over 1 fathom of water; 37 specimens.

Station 40m: Bernard harbour, Northwest Territories, June 25, 1915, from stomach of *Xema sabini* (J. Sabine); 2 specimens.

Station 41g: Bernard harbour, Northwest Territories (outer harbour), August 1, 1915, surface; 2 specimens.

Station 41u: Bernard harbour, Northwest Territories, end of August, 1915, from stomach of *Salvelinus malma* Walb.; 44 specimens.

Station 42h: Bay at Bernard harbour, Northwest Territories, September 22, 1915, beach water; 12 specimens.

Station 46g: Dolphin and Union strait (Bernard harbour), Northwest Territories, May 4, 1916, from Eskimos; 20 specimens.

Station 61a: South of Armstrong point, Victoria island, Prince of Wales strait, Northwest Territories, October, 1915, J. Hadley collector; 200 specimens.

Distribution.—A very abundant species found throughout the Arctic ocean.

10. *Pseudalibrotus nanseni* G. O. Sars.

1900. *Pseudalibrotus nanseni* G. O. Sars, in NANSEN, Norwegian North Polar Exp., vol. I, No. 5, p. 26, pl. 4, 5.

Station 57a: Cape Smyth (point Barrow) Alaska, August 8, 1916, pelagic, over 1 fathom of water; 1 specimen.

The specimens described by Sars were collected by the Norwegian North Polar Expedition in 1894 and 1896 north of the New Siberian islands, and farther to

the west. The discovery of this specimen at point Barrow extends the range of this species considerably to the east.

Juvenile specimens of some form of *Pseudalibrotus* were obtained at Station 40r, 50b (both in Dolphin and Union strait), and 56a, (Harrison bay, Alaska), but any specific identification of such specimens would be very doubtful.

11. *Socarnes bidenticulatus* (Bate).

1835. *Gammarus nugax* J. C. Ross, in Ross's Second voyage, Appendix, p. 87.
 1906. *Socarnes bidenticulatus* STEBBING, Tierreich, Amph. I, p. 56, and synonymy.
 1909. *Socarnes bidenticulatus* BRÜGGEN, Mém. Acad. Imp. Sci. de St. Pétersbourg, sér. 8, vol 18, No. 16, p. 12.
 1912. *Socarnes bidenticulatus* STEPHENSEN, Report on the Malacost., Pycnogonida, etc., collected by the Danmark Exped. to Northeast Greenland, p. 527.
 1913. *Socarnes bidenticulatus* STEPHENSEN, Account Crust. etc. collected by Dr. V. Nordmann in Summer 1911 from West Greenland, p. 65.

Station 42u: Bernard harbour, Northwest Territories, October 22, 1915, from stomach of *Erignathus barbatus* (Erxleben); 5 specimens.

Distribution.—Arctic ocean, east and west coast of Greenland.

12. *Tmetonyx gulosus* (Krøyer).

1845. *Anonyx gulosus* KRØYER, Naturh. Tidsskr., ser. 2, vol. I, p. 611.
 1906. *Tmetonyx cicada* STEBBING, Tierreich, Amph. I, p. 74, and synonymy.
 1909. *Tmetonyx cicada* BRÜGGEN, Mém. Acad. Imp. Sci. St. Pétersbourg, sér. 8, vol. 18, No. 16, p. 13.
 1911. *Tmetonyx gulosus* STAPPERS, Crust. Malacost., Campagne Arctique de 1907 du Duc d'Orléans, p. 11, and synonymy.

Station 29f: Lat. 70° 13' N., long. 140° 50' W., April 4, 1914, from stomach of *Phoca hispida* Schreber, over about 30 fathoms; 4 specimens.

Station 42u: Bernard harbour, Northwest Territories, October 22, 1915, from stomach of *Erignathus barbatus* (Erxleben); 1 specimen.

Station 43a: Dolphin and Union strait (off Cockburn point), Northwest Territories, September 13, 1915, over about 50 fathoms of water, mud with pebbles, but no algæ; 2 specimens.

Station 62a: Liddon gulf, Melville island, Northwest Territories, July, 1916, A. Castel collector; 2 specimens.

Distribution.—Arctic ocean, north Atlantic, North sea, France.

13. *Katius obesus* Chevreux.

1905. *Katius obesus* CHEVREUX, Description d'un Amphipode (*Katius obesus* nov. gen. et sp.) Bull. Mus. Océanogr. Monaco, No. 35, 1915 (with figs).
 1906. *Katius obesus* TATTERSALL, Fisheries, Ireland, Sci. Invest. 1905, IV. 1906, p. 29.
 1912. *Katius obesus* STEPHENSEN, Vidensk. Meddel. fra den Naturh. Foren., Bd. 64, p. 89.

Station 8a: Lat. 55° 13' N., long. 140° 21' W., June 26, 1913, surface; 1 specimen.

This species is represented by a single badly preserved specimen which differs only slightly from Chevreux's figure. His specimen was 12 mm. long, while those obtained on the *Tjalfe* Expedition were 25 mm. The present specimen measures about 5 mm. which indicate that it is quite immature. The presence of a double row of calceoli on each antenna would seem to indicate a male. Chevreux's specimen apparently did not have the calceoli, but he was not certain that it was a male. The eyes in our specimen are not present, as nearly all of the internal organs have disappeared, leaving the interior entirely empty. The 3rd uropods have the rami much narrower and the inner ramus proportionately shorter than in Chevreux's figure. The type specimen was obtained off the Azores, in water 0-3000 meters depth; the specimen obtained by the *Helga* came from the west coast of Ireland, 1,200 fathoms, and the specimens collected by the *Tjalfe* Expedition were from the southern coast of Green-

land, in water 500-2000 meters depth. The discovery of this specimen in the North Pacific upon the surface greatly extends its range, both geographically and bathymetrically.

Family **STEGOCEPHALIDÆ**.

14. **Stegocephalus ampulla** (Phipps).

1774. *Cancer ampulla* PHIPPS, Voy. North Pole, p. 191, pl. 12, f. 3.
 1906. *Phippsia ampulla* STEBBING, Tierreich, Amph. I, p. 89, and synonymy.
 1909. *Stegocephalus ampulla* BRÜGGEN, Mém. Acad. Imp. Sci. St. Pétersbourg, sér. 8, vol. 18, No. 16, p. 14; pl. I, fig. 1; pl. III, figs. 11-19.

Station 42u: Bernard harbour, Northwest Territories, October 22, 1915, from stomach of *Erignathus barbatus* (Erxleben); 1 specimen.
Distribution.—Arctic ocean.

15. **Stegocephalus inflatus** Krøyer.

1842. *Stegocephalus inflatus* KRØYER, Naturh. Tidsskr., vol. 4, p. 150.
 1906. *Stegocephalus inflatus* STEBBING, Tierreich, Amph. I, p. 91, and synonymy.
 1909. *Stegocephalus inflatus* BRÜGGEN, Mém. Acad. Imp. Sci. St. Pétersbourg, sér. 8, vol. 18, No. 16, p. 14.
 1911. *Stegocephalus inflatus* STAPPERS, Crust. Malacost., Campagne Arctique de 1907 du Duc d'Orléans, p. 28.
 1912. *Stegocephalus inflatus* STEPHENSEN, Vidensk. Meddel. fra den Naturh. Foren., Bd. 64, p. 89.
 1912. *Stegocephalus inflatus* STEPHENSEN, Meddel. om Grønland, vol. XLV, p. 532.
 1913. *Stegocephalus inflatus* STEPHENSEN, Meddel. om Grønland, vol. LI, p. 66.
 1916. *Stegocephalus inflatus* STEPHENSEN, Meddel. om Grønland, vol. LIII, p. 286.

Station 29f: Lat. 70° 13' N., long. 140° 50' W., April 4, 1914, from stomach of *Phoca hispida* Schreber, about 30 fathoms; 6 specimens.

Station 42u: Bernard harbour, Northwest Territories, October 22, 1915, from stomach of *Erignathus barbatus* (Erxleben); 10 specimens.

Station 43a: Dolphin and Union strait (off Cockburn point), Northwest Territories, September 13, 1915. 100 meters; sandy mud with pebbles but no algae; 1 specimen.

Distribution.—Arctic Ocean; north Atlantic (west-Norway, Shetland Isles, Nova Scotia, Labrador, Massachusetts); north Pacific (Japan, sea of Okhotsk); Bering sea.

Family **AMPELISCIDÆ**.

16. **Ampelisca eschrichtii** Krøyer.

1842. *Ampelisca eschrichtii* KRØYER, Naturh. Tidsskr., vol. 4, p. 155.
 1906. *Ampelisca eschrichtii* STEBBING, Tierreich, Amph. I, p. 100, and synonymy.
 1906. *Ampelisca eschrichtii* CHEVREUX, Amph. Exped. Antaret. franc., p. 20.
 1909. *Ampelisca eschrichtii* BRÜGGEN, Mém. Acad. Imp. Sci. St. Pétersbourg, sér. 8, vol. 18, No. 16, p. 16.
 1911. *Ampelisca eschrichtii* STAPPERS, Crust. Malacost. Campagne Arctique de 1907 du Duc d'Orléans, p. 19, pl. I, fig. 1, 9, 14-16.
 1917. *Ampelisca eschrichtii* CHILTON, Journal of Zool. Research, May, 1917, vol. II, No. 2, p. 87.

Station 43a: Dolphin and Union strait (off Cockburn point), Northwest Territories, September 13, 1915; about 50 fathoms, mud with pebbles but no algae; 1 specimen.

Distribution.—Arctic ocean, north Atlantic, North and South Pacific, Bay of Biscoe, Anvers Island (about Lat. 64° S. Long. 64° W.)

17. **Haploops tubicola** Lilljeborg.

1855. *Haploops tubicola* LILLJEBORG, Öfv. Ak. Förh., vol. 12, p. 135, 136.
 1906. *Haploops tubicola* STEBBING, Tierreich, Amph. I, p. 117, and synonymy.
 1908. *Haploops tubicola* HOLMES, Proc. U.S.N.M., vol. XXXV, p. 518.

1909. *Haploops tubicola* BRÜGGEN, Mém. Acad. Imp. Sci. St. Pétersbourg, sér. 8, vol. 18, No. 16, p. 19.
 1913. *Haploops tubicola* STEPHENSEN, Meddel. om Grønland, vol. LI, p. 67.
 1916. *Haploops tubicola* STEPHENSEN, Meddel. om Grønland, vol. LIII, p. 294.

Station 43b: Dolphin and Union strait (off Staphylton bay), Northwest Territories, September 14, 1915, 25-30 fathoms, sandy mud with pebbles but no algæ; 1 specimen.

Distribution.—Arctic ocean, North Atlantic, North Pacific.

Family HAUSTORIIDÆ.

18. *Pontoporeia affinis* Lindström.

1855. *Pontoporeia affinis* LINDSTRÖM, Öfv. Ak. Förh., vol. 12, p. 63.
 1906. *Pontoporeia affinis* STEBBING, Tierreich, Amph. I, p. 128, and synonymy.
 1909. *Pontoporeia affinis* BRÜGGEN, Mém. Acad. Imp. Sci. St. Pétersbourg, sér. 8, vol. 18, No. 16, p. 21.

Station 27o: Collinson point, Alaska, September 20, 1913, pelagic, over about 1 foot of water (9-inch ice); 2 specimens.

Station 28o: Collinson point, Alaska, July 8, 1914, from stomach of *Salvelinus malma* Walb.; 5 specimens.

Distribution.—Fresh-water lakes (Norway, Sweden, Russia, North America); Baltic, Kattegat, Kara sea, North Atlantic (France).

19. *Pontoporeia femorata* Krøyer.

1842. *Pontoporeia femorata* KRØYER, Naturh. Tidsskr., vol. 4, p. 153.
 1906. *Pontoporeia femorata* STEBBING, Tierreich, Amph. I, p. 128, and synonymy.
 1909. *Pontoporeia femorata* BRÜGGEN, Mém. Acad. Imp. Sci. St. Pétersbourg, sér. 8, vol. 18, No. 16, p. 20.

Station 37j: Bernard harbour, Northwest Territories, September 1, 1914, pelagic, over about 2 fathoms of water; 1 specimen.

Station 41u: Bernard harbour, Northwest Territories, end of August, 1915, from stomach of *Salvelinus malma* Walb.; 2 specimens.

Distribution.—Arctic ocean, North Atlantic, Baltic.

20. *Priscillina armata* (Boeck).

1861. *Pontoporeia armata* BOECK, Forh. Skand. Naturf., Møde 8, p. 648.
 1906. *Priscillina armata* STEBBING, Tierreich, Amph. I, p. 126, and synonymy.
 1913. *Priscilla armata* STEPHENSEN, Meddel. om Grønland, vol. XXII, p. 129.

Station 48b: Banks peninsula, Bathurst inlet, Northwest Territories, May 18, 1916, from stomach of *Gadus* sp.; 1 specimen.

Distribution.—Arctic ocean, North Atlantic, Greenland, West Norway?

Family ACANTHONOTOZOMATIDÆ.

21. *Acanthonotozoma inflatum* (Krøyer).

1842. *Acanthonotus inflatus* KRØYER, Naturh. Tidsskr., vol. 4, p. 161.
 1906. *Acanthonotozoma inflatum* STEBBING, Tierreich, Amph. I, p. 219 and synonymy.
 1909. *Acanthonotozoma inflatum* BRÜGGEN, Mém. Acad. Imp. Sci. St. Pétersbourg, sér. 8, vol. 18, No. 16, p. 23.

Station 28d: Collinson point, Alaska, October 18, 1913, about 1 fathom; 1 specimen.

Station 28e: Collinson point, Alaska, October 21, 1913, pelagic, over about 1 fathom; 1 specimen.

Distribution.—Arctic ocean, Greenland.

Family OEDICEROTIDÆ.

22. *Acanthostephia malmgreni* (Goës).

1806. *Amphilnotus malmgreni* GOËS, Ofv. Ak. Förh., vol. 22, p. 526, pl. 39, f. 17.
 1906. *Acanthostephia malmgreni* STEBBING, Tierreich, Amph. I, p. 254, and synonymy.
 1909. *Acanthostephia malmgreni* BRÜGGEN, Mém. Acad. Imp. Sci. St. Pétersbourg, sér. 8, vol. 18, No. 16, p. 25.

Station 43b: Dolphin and Union strait, Northwest Territories (off Stapyllton bay), September 14, 1915, 25-30 fathoms; sandy mud with pebbles, but no algæ; 1 specimen.

Distribution.—Arctic ocean, Greenland.

23. *Acanthostephia pulchra* Miers.

1881. *Acanthostephia pulchra* MIERS, Ann. Nat. Hist., ser. 5, vol. 7, p. 47, pl. 7, f. 1, 2.
 1909. *Acanthostephia pulchra* STEBBING, Tierreich, Amph. I, p. 254, and synonymy.

Station 25a: Off Cooper island (point Barrow), Alaska, pelagic, over 0-2 fathoms, August 27, 1913; 5 specimens.

Station 27s: Collinson point, Alaska, October 3, 1913, about 3 fathoms; 1 specimen.

Station 28o: Collinson point, Alaska, July 8, 1914, from stomach of *Salvelinus malma* Walb.; 13 specimens.

Station 41c: Bernard harbour, Northwest Territories, July 28, 1915, about 5 fathoms, sandy mud with algæ; 3 specimens.

Station 41u: Bernard harbour, Northwest Territories, end of August, 1915, from stomach of *Salvelinus malma* Walb.; 2 specimens.

Station 59a: Off Cape Kellett, Banks island, Northwest Territories, September 7, 1914, 5-6 fathoms, sand with algæ; G. H. Wilkins, collector; 2 specimens.

Color.—Entire animal light grayish vinaceous with the body segments each transversely barred with dark purple-drab, joints of peduncles of antennae transversely barred with dark purple-drab.

Distribution.—Arctic ocean (Franz Josef land, Siberia).

24. *Aceroides latipes* (G. O. Sars).

1882. *Halicreion latipes* SARS, Forh. Selsk. Christian., nr. 18, p. 97, t. 4, f. 10.
 1906. *Aceroides latipes* STEBBING, Tierreich, Amph. I, p. 255, and synonymy.
 1909. *Aceroides latipes* BRÜGGEN, Mém. Acad. Imp. Sci. St. Pétersbourg, sér. 8, vol. 18, No. 16, p. 24.

Station 27s: Collinson point, Alaska, October 3, 1913, about 3 fathoms, mud with algæ; 1 specimen.

Station 28o: Collinson point, Alaska, July 8, 1914, from stomach of *Salvelinus malma* Walb.; 1 specimen.

Distribution.—Arctic ocean, Greenland, Norway.

25. *Arrhis phyllonyx* (M. Sars).

1858. *Leucothoe phyllonyx* M. SARS, Forh. Selsk. Christian., p. 148.
 1906. *Arrhis phyllonyx* STEBBING, Tierreich, Amph. I, p. 248, and synonymy.
 1909. *Arrhis phyllonyx* BRÜGGEN, Mém. Acad. Imp. St. Pétersbourg, sér. 8, vol. 18, No. 16, p. 25.

Station 43b: Dolphin and Union strait (off Stapyllton bay), Northwest Territories, September 14, 1915, 25-30 fathoms, sandy mud with pebbles, but no algæ; 1 specimen.

Distribution.—Arctic ocean, North Atlantic, North sea, Greenland, Iceland, Norway.

26. *Monoculodes longirostris* (Goës).

1886. *Oediceros longirostris* GOËS, Öfv. Ak. Förh., vol. 22, p. 526, pl. 39, f. 20.

1906. *Monoculodes longirostris* STEBBING, Tierreich, Amph. I, p. 260, and synonymy.

Station 37j: Bernard harbour, Northwest Territories, September 1, 1914, pelagic, over 2 fathoms water; 1 specimen.

Distribution.—Arctic ocean (Spitzbergen, Finmark, Tromsø), Kattegat.

27. *Monoculodes schneideri* G. O. Sars.

1895. *Monoculodes schneideri* SARS, Crust. Norway., vol. I, p. 692, pl. VI, f. 1.

1906. *Monoculodes schneideri* STEBBING, Tierreich, Amph. I, p. 263.

Station 57a: Cape Smyth (point Barrow), Alaska, August 8, 1916, pelagic, over 1 fathom water; 2 specimens.

Distribution.—Arctic ocean (Tromsø).

One immature specimen of some form of *Monoculodes* was obtained at station 41n (Bernard harbour).

28. *Monoculopsis longicornis* (Boeck).

1871. *Monoculodes longicornis* BOECK, Forh. Selsk. Christian., 1870, p. 165.

1906. *Monoculopsis longicornis* STEBBING, Tierreich, Amph. I, p. 258, and synonymy.

1912. *Monoculopsis longicornis* STEPHENSEN, Meddel. om Grønland, vol. XLV, p. 505.

Station 41n: Bernard harbour, Northwest Territories, (inner harbour), August 9, 1915, surface; 2 specimens.

Distribution.—Arctic ocean, North Atlantic, North sea, Greenland.

29. *Paroediceros lynceus* (M. Sars).

1858. *Oediceros lynceus* SARS, Forh. Selsk. Christian., p. 143.

1906. *Paroediceros lynceus* STEBBING, Tierreich, Amph. I, p. 246, and synonymy.

1909. *Paroediceros lynceus* BRÜGGEN, Mém. Acad. Imp. Sci. St. Pétersbourg, sér. 8, vol. 18, No. 16, p. 23.

1911. *Paroediceros lynceus* STAPPERS, Crust. Malacost. Campagne Arctique de 1907, du Duc d'Orléans, p. 72.

1912. *Paroediceros lynceus* STEPHENSEN, Meddel. om Grønland, vol. XLV, p. 532.

1913. *Paroediceros lynceus* STEPHENSEN, Meddel. om Grønland, vol. LI, p. 66.

1916. *Paroediceros lynceus* STEPHENSEN, Meddel. om Grønland, vol. LIII, p. 287.

Station 27s: Collinson point, Alaska, October 3, 1913, 3 fathoms, mud with algae; 1 specimen.

Station 28o: Collinson point, Alaska, July 8, 1914, from stomach of *Salvelinus malma* Walb; 2 specimens.

Distribution.—Arctic ocean (Greenland, Iceland, Spitzbergen, Murman coast, Siberian Polar sea, Labrador, Finland).

Family CALLIOPIIDÆ.

30. *Apherusa glacialis* (H. J. Hansen).

1887. *Apherusa glacialis* HANSEN, Vid. Meddel. ser. 4, vol. 9, p. 137, pl. 5, f. 6-6c.

1906. *Apherusa glacialis* STEBBING, Tierreich, Amph. I, p. 307, and synonymy.

1909. *Apherusa glacialis* BRÜGGEN, Mém. Acad. Imp. Sci. St. Pétersbourg, sér. 8, vol. 18, No. 16, p. 28.

1911. *Apherusa glacialis* STAPPERS, Crust. Malacost., Campagne Arctique de 1907 du Duc d'Orléans, p. 61.

1912. *Apherusa glacialis* STEPHENSEN, Vidensk. Meddel. fra den Naturh. Foren., København, Bd. 64, p. 96.

1912. *Apherusa glacialis* STEPHENSEN, Meddel. om Grønland, vol. XLV, p. 537.

1916. *Apherusa glacialis* STEPHENSEN, Meddel. om Grønland, vol. LIII, p. 289.

Station 25b, c: Off Cooper island, (point Barrow), Alaska, August 27, 1913, surface; 1 specimen.

Station 41r: Beach at Bernard harbour, Northwest Territories, August 14, 1915; 2 specimens.

Station 56a: Harrison bay, Alaska, August 6, 1916, surface; 1 specimen.

Station 57a: Cape Smyth (point Barrow), Alaska, August 8, 1916, pelagic, over 1 fathom of water; 3 specimens.

Distribution.—Arctic ocean, Greenland.

31. *Apherusa megalops* (Buchholz).

1874. *Paramphithoe megalops* BUCHHOLZ, Zweite Deutsche Nordpolarfahrt, vol. 2, p. 369, pl. 12.

1906. *Halirages megalops* STEBBING, Tierreich, Amph. I, p. 293, and synonymy.

1912. *Amphithopsis megalops* STEPHENSEN, Meddel. om Grønland, vol. XLV, p. 584.

1913. *Amphithopsis megalops* STEPHENSEN, Meddel. om Grønland, vol. XXII, p. 176.

1916. *Amphithopsis megalops* STEPHENSEN, Meddel. om Grønland, vol. LIII, p. 290.

Station 37j: Bernard harbour, Northwest Territories, September 1, 1914, pelagic, over 2 fathoms of water; 2 specimens.

Station 41: Bernard harbour, Northwest Territories (outer harbour), July 20, 1915, 5 fathoms; 1 specimen.

This is not the *Apherusa megalops* described by G. O. Sars in 1882 from Norway, but the species described by R. BUCHHOLZ in 1874 as *Paramphithoe megalops* from Northeast Greenland. Heretofore this species has not been recorded outside of Greenland.

32. *Calliopius laevisculus* (Krøyer).

1838. *Amphithoe laeviscula* KRØYER, Danske Selsk. Afh., vol. 7, p. 281, pl. 3, f. 13 a-h.

1906. *Calliopius laevisculus* STEBBING, Tierreich, Amph. I, p. 296, and synonymy.

1912. *Calliopius laevisculus* STEPHENSEN, Meddel. om Grønland, vol. XLV, p. 597, 617.

1913. *Calliopius laevisculus* STEPHENSEN, Meddel. om Grønland, vol. XXII, p. 179.

Station 7a: Lat. 55° 42' N., long. 136° 20' W., June 25, 1913, surface (among floating algæ); 3 specimens.

Station 13a, b, c: Lat. 54° 30' N., long. 159° 42' W., July 1, 1913, surface; 1 specimen.

Station 13g, h: Lat. 54° 30' N., long. 159° 42' W., July 1, 1913, surface; 2 specimens.

Station 14: Lat. 54° 23' N., long. 164° 45' W., July 2, 1913, surface; 1 specimen.

Distribution.—Arctic ocean, North Atlantic, North sea, Greenland, Labrador, Norway, British Isles, North Pacific.

33. *Calliopius rathkii* (Zaddach).

1844. *Amphithoe rathkii* ZADDACH, Synops. Crust. Pruss., p. 6.

1906. *Calliopius rathkii* STEBBING, Tierreich, Amph. I, p. 296, and synonymy.

1912. *Calliopius rathkei* STEPHENSEN, Meddel. om Grønland, vol. XLV, p. 597.

1913. *Calliopius rathkei* STEPHENSEN, Meddel. om Grønland, vol. XXII, p. 180.

1916. *Calliopius rathkei* STEPHENSEN, Meddel. om Grønland, vol. LIII, p. 292.

Station 41s: Bernard harbour, Northwest Territories (inner harbour), August 24, 1915, surface; 1 specimen.

Distribution.—Arctic ocean, North Atlantic, North sea, and Skagerrak, Norway, France, Great Britain.

34. *Halirages nilssoni* Ohlin.

1895. *Halirages nilssoni* A. OHLIN, Acta Univ. Lund., vol. 31, No. 6, p. 44, pl. f. 1-6.

Station 41: Bernard harbour, Northwest Territories (outer harbour), July 20, 1915, 5 fathoms, sandy mud with algæ; 3 specimens.

This species was first obtained from Baffin bay in 1894 by the Peary Auxiliary Expedition. The present specimens from Bernard harbour make the second record, and extend the range considerably to the west.

Family ATYLIDÆ.

35. *Atylus carinatus* (Fabricius).

1793. *Gammarus carinatus* FABRICIUS, Ent. Syst., vol. 2, p. 515.
 1906. *Atylus carinatus* STEBBING, Tierreich, Amph. I, p. 328, and synonymy.
 1909. *Atylus carinatus* BRÜGGEN, Mém. Acad. Imp. Sci. St. Pétersbourg, sér. 8, vol. 18, No. 16, p. 32.
 1912. *Atylus carinatus* STEPHENSEN, Meddel. om Grønland, vol. XLV, p. 540, 605.
 1913. *Atylus carinatus* STEPHENSEN, Meddel. om Grønland, vol. XXII, p. 171.

Station 20b, Grantley harbour (port Clarence), Alaska, July 30, 1913, 2-3 fathoms, sandy mud with many algæ; 5 specimens.

Station 20g: Port Clarence, Alaska, August 4, 1913, 2-3 fathoms, mud with many algæ, 2 specimens.

Station 27r: Collinson point, Alaska, October 2, 1913, pelagic, over 1 fathom water; 3 specimens.

Station 27s: Collinson point, Alaska, October 3, 1913, 3 fathoms, mud with algæ; 5 specimens.

Station 28d: Collinson point, Alaska, October 18, 1913, pelagic; 2 specimens.

Station 28e: Collinson point, Alaska, October 21, 1913, pelagic, over 1 fathom water; 1 specimen.

Station 28f: Collinson point, Alaska, October 25, 1913, 1½ fathoms, sand with algæ; 2 specimens.

Station 40u: Bernard harbour, Northwest Territories, July 6-8, 1915, from stomach of *Erignathus barbatus* (Erxleben); 1 specimen.

Station 41: Bernard harbour, Northwest Territories (outer harbour), July 20, 1915, 5 fathoms, sandy mud with many algæ; 21 specimens.

Color.—Entire animal tawny-olive with a sepia spot on the lower part of the body segments, carina marked in front with sepia, second joint of peduncle of the first and second antennae darker than the rest.

Distribution.—Arctic ocean (widely distributed).

Family EUSIRIDÆ.

36. *Rhachotropis aculeata* (Lepechin).

1780. *Oniscus aculeata* LEPECHIN, Acta Ac. Petrop., 1778i, p. 247, pl. 8, f. 1.
 1906. *Rhachotropis aculeata* STEBBING, Tierreich, Amph. I, p. 348, and synonymy.
 1909. *Rhachotropis aculeata* BRÜGGEN, Mém. Acad. Imp. Sci. St. Pétersbourg, sér. 8, vol. 18, No. 16, p. 34.
 1916. *Rhachotropis aculeata* STEPHENSEN, Meddel. om Grønland, vol. LIII, p. 292.

Station 29f: Lat. 70° 13' N., long. 140° 50' W., April 4, 1914, from stomach of *Phoca hispida* (Schreber), 30 fathoms; 1 specimen.

Distribution.—Arctic ocean (widely distributed, circumpolar).

A fragment of some species of *Rhachotropis* was collected at station 28f, Collinson point, Alaska.

37. *Rozinante fragilis* (Goës).

1866. *Paramphithoe fragilis* GOËS, Öfv. Ak. Förh., vol. 22, p. 524, pl. 39, f. 16.
 1906. *Rozinante fragilis* STEBBING, Tierreich, Amph. I, p. 354, and synonymy.
 1909. *Rozinante fragilis* BRÜGGEN, Mém. Acad. Imp. Sci. St. Pétersbourg, sér. 8, vol. 18, No. 16, p. 35.
 1911. *Rozinante fragilis* STAPPERS, Crust. Malacost., Campagne Arctique de 1907, du Duc d'Orléans, p. 56, pl. III, f. 1-4.

Station 27s: Collinson point, Alaska, October 3, 1913, 3 fathoms, mud with algæ. (Bottom and pelagic); 2 specimens.

Station 28e: Collinson point, Alaska, October 21, 1913, pelagic, over 1 fathom of water; 2 specimens.

Distribution.—Arctic ocean (Greenland, Spitzbergen, Barents sea).

Family PONTOGENEIIDÆ.

38. *Pontogeneia inermis* (Krøyer).

1838. *Amphithoe inermis* KRØYER, Danske Sebk. Aft., vol. 7, p. 275, pl. 3, f. 11 a-g.
 1838. *Amphithoe crenulata* KRØYER, *ibid.*, p. 278, pl. 3, f. 12 a-g.
 1906. *Pontogeneia inermis* STEBBING, Tierreich, Amph. I, p. 359, and synonymy.
 1912. *Pontogeneia inermis* STEPHENSEN, Meddel. om Grønland, vol. XLV, p. 539, 606.
 1913. *Pontogeneia inermis* PEARSE, Proc. U.S.N.M., vol. 45, p. 573.
 1913. *Pontogeneia inermis* STEPHENSEN, Meddel. om Grønland, vol. XXII, p. 173.
 1916. *Pontogeneia inermis* STEPHENSEN, Meddel. om Grønland, vol. LIII, p. 289.

Station 37j: Bernard harbour, Northwest Territories, September 1, 1914, pelagic, over 2 fathoms water; 1 specimen.

Distribution.—Arctic ocean, North Atlantic, North Pacific, North sea, Greenland, West Norway, Siberia.




Figure 3. *Rozinante fragilis* (Goëss).

1 Peraeopod 2. 2 Gnathopod 2.

Family GAMMARIDÆ.

39. *Gammaracanthus loricatus* (Sabine).

- 1821 and 24. *Gammarus loricatus* E. SABINE, in W. E. Parry, J. Voy., Suppl., p. 53, pl. I, f. 7; p. 231, pl. I, f. 7.
 1906. *Gammaracanthus loricatus* STEBBING, Tierreich, Amph. I, p. 508, and synonymy.
 1909. *Gammaracanthus loricatus* BRÜGGEN, Mém. Acad. Imp. Sci. St. Pétersbourg, sér. 8, vol. 18, No. 16, p. 35.
 1912. *Gammaracanthus loricatus* STEPHENSEN, Meddel. om Grønland, vol. XLV, p. 503, 543, 589.
 1913. *Gammaracanthus loricatus* STEPHENSEN, Meddel. om Grønland, vol. XXII, p. 197.

Station 28c: Collinson point, Alaska, October 14, 1913, sandy mud with scattered algæ, 1 fathom; 1 specimen.

Station 29f: Lat. 70° 13' N., long. 140° 50' W., April 4, 1914, water depth 30 fathoms, from stomach of *Phoca hispida* Schreber; 2 specimens.

Station 28o: Collinson point, Alaska, July 8, 1914, from stomach of *Salvelinus malma* Walb.; 2 specimens.

Station 37j: Bernard harbour, Northwest Territories, September 1, 1914, pelagic, over 2 fathoms water; 4 specimens.

Station 37r, u: Bernard harbour, Northwest Territories, October 16-20, 1914, pelagic, over 1 fathom of water; 1 specimen.

Station 40u: Bernard harbour, Northwest Territories, July 6-8, 1915, from stomach of *Erignathus barbatus* (Erxleben); 1 specimen.

Station 41: Bernard harbour, Northwest Territories (outer harbour), July 20, 1915, sandy mud with many algæ, 5 fathoms; 3 specimens.

Distribution.—Arctic Ocean (Greenland, Spitzbergen, Franz Josef land, Nova Zembla, Siberia.)

40. *Gammarus limnæus* Smith.

1871. *Gammarus locustis* S. I. SMITH, Amer. J. Sci., ser. 3, vol. 2, p. 453.
 1874. *Gammarus limnæus*, S. I. SMITH, 7th Rept. U.S. Geol. Survey, p. 609; Rept. U.S. Fish Com. 1872-73, p. 651.
 1874. *Gammarus robustus* S. I. SMITH, 7th Rept. U.S. Geol. Survey, p. 610.
 1907. *Gammarus limnæus* A. L. WECKEL, Proc. U.S.N.M., vol. XXXII, p. 42, f. 9.

Station 28h: Warm creek, tributary to Sadlerochit river, about 25 miles inland from Camden bay, Alaska, November 6-7, 1913: 30 specimens.

Station 40g: Bernard harbour, Northwest Territories, lake or creek, from stomach of *Cristivomer namaycush* Walb., June 28, 1915: 17 specimens.

Station 40n¹: Lake, inland at Bernard harbour, Northwest Territories, June 23, 1915, pelagic, over 4 meters of water, ice 2 meters; 11 specimens.

Station 40n²: Pond at Bernard harbour, Northwest Territories, July 16, 1915, pelagic; 3 specimens.

Station 42r: Lake at Bernard harbour, Northwest Territories, October 2, 1915, from stomach of *Salvelinus marstoni* Garm.; 5 specimens.

Station 42v: Lake at Bernard harbour, Northwest Territories, December, 1915, from stomach of *Salvelinus marstoni* Garm.; 7 specimens.

Station 50g: Lake at Bernard harbour, Northwest Territories, April, 1906, from stomach of *Cristivomer namaycush* Walb.; 4 specimens.

Station 54h: Pond on north side of east end of Herschel island, Yukon Territory, August 1, 1916; 28 specimens.

This Amphipod has been found throughout the northeastern and western parts of the United States, and these records now from Bernard harbour and Alaska indicate that it inhabits the fresh-waters of the United States, Canada, and Alaska.

41. *Gammarus locusta* (Linné).

1758. *Cancer locusta* LINNÉ, Syst. Nat., ed. 10, p. 634.
 1906. *Gammarus locusta* STEBBING, Tierreich, Amph. I, p. 476, and synonymy.
 1909. *Gammarus locusta* BRÜGGEN, Mém. Acad. Imp. Sci. St. Pétersbourg, sér. 8, vol. 18, No. 16, p. 35.
 1911. *Gammarus locusta* STAPPERS, Crust. Malacost., Campagne Arctique de 1907 du Duc d'Orléans, p. 68.
 1913. *Gammarus locusta* PEARSE, Proc. U.S.N.M., vol. 45, p. 571.
 1916. *Gammarus locusta* STEPHENSEN, Meddel. om Grønland, vol. LIII, p. 293.

Station 20b, c: Grantley harbour (port Clarence), Alaska, July 30, 1913, 2-3 fathoms, sandy mud with many algæ; 2 specimens.

Station 20h: Port Clarence, Alaska, August 4, 1913, 2-3 fathoms, surface, attached to floating algæ; 3 specimens.

Station 27a: Collinson point, Alaska, September 5, 1913, sandy mud with scattered algæ, from stomach of *Cottus quadricornis* L.; 2 specimens.

Station 27c: Lagoon-bay at Collinson point, Alaska, September 15, 1913, 0-1 foot water; 1 specimen.

Station 27h: Lagoon-bay at Collinson point, Alaska, 0-1 foot water, September 18, 1913; 15 specimens.

Station 27o: Collinson point, Alaska, September 20, 1913, pelagic, over 1 foot of water (9-inch ice); 1 specimen.

Station 28c: Collinson point, Alaska, October 14, 1913, 1 fathom, sandy mud with scattered algæ; 35 specimens.

Station 28l: Beach at Collinson point, Alaska, June, 1914; 2 specimens.

Station 28o: Collinson point, Alaska, from stomach of *Salvelinus malma* Walb., July 8, 1914; 14 specimens.

Station 28r: Bay at Collinson point, Alaska, July 24, 1914; from stomach of *Collus quadricornis* L. 2 fragments.

Station 37a: Bernard harbour, Northwest Territories, August 24, 1914, from stomach of *Erignathus barbatus* (Erxleben); 5 specimens.

Station 37j: Bernard harbour, Northwest Territories, September 1, 1914, pelagic, over 2 fathoms water; 1 specimen.

Station 37r, u: Bernard harbour, Northwest Territories, October 16-20, 1914, pelagic, over 1 fathom water; 6 specimens.

Station 40d: Dolphin and Union strait (Bernard harbour), Northwest Territories, June 8, 1915. Pelagic over 9 fathoms; 1 specimen.

Station 40h: Dolphin and Union strait (Bernard harbour), Northwest Territories, June 25, 1915, pelagic, over 2½ fathoms water; 5 specimens.

Station 40m: Bernard harbour, Northwest Territories, June 25, 1915, from stomach of *Xema sabini* (J. Sabine); 1 specimen.

Station 40p: Bernard harbour, Northwest Territories, July 1, 1915, pelagic, in littoral region; 20 specimens.

Station 40u: Bernard harbour, Northwest Territories, July 6-8, 1915, from stomach of *Erignathus barbatus* (Erxleben), F.; 4 specimens.

Station 48b: Banks peninsula, Bathurst inlet, Northwest Territories, May 18, 1916, from stomach of *Gadus* sp.; 10 specimens.

Station 50d: Young point, Northwest Territories (Dolphin and Union strait), July 21, 1916, among loose algæ, in littoral region; 6 specimens.

Color.—Entire animal maize yellow, eyes berlin blue, legs and antennae banded with light russet-vinaceous, each of the body segments with a dark, transverse dorsal band, the last two thoracic and the abdominal segments marked on their lower portions with scarlet.

Distribution.—Arctic ocean, North Atlantic, North Pacific.

Immature species of *Gammarus* were obtained at station 40r, 41g, (both Dolphin and Union strait), and 56a (Harrison Bay, Alaska).

42. *Synurella johanseni*, new species.

Type specimen: Catalogue No. 1380; paratypes, Nos. 1381-3, Victoria Memorial Museum, Ottawa, Canada.

Station 20i: Pond in the tundra at Teller, Alaska, August 6, 1913; 21 specimens.

Station 20i: Brackish pond at Teller, Alaska, August 3, 1913; 1 specimen (juvenile).

Description.—Body rather stout, not much compressed. Head with front edge broadly rounded. Eyes rather small, irregularly oval, composed of few elements, brownish black.

Pleon segments 1 and 2 with posterior lateral corners slightly produced; segment 3, with posterior lateral corners evenly rounded or with a very slight production of the lower posterior margin. Posterior edge of pleon segment 3 with shallow notch bearing a minute seta just above the lower margin.

Antenna 1 is about half as long as the body; the joints of the peduncle becoming consecutively shorter; 1st joint much the stoutest; flagellum of about

12 joints; accessory flagellum 2-jointed, but not as long as the 1st joint of primary.

Antenna 2 with 4th joint of peduncle longest; flagellum 6-jointed, and about as long as the 4th joint of peduncle.

Side-plates 1-3 about as deep as their segments, with sides parallel and lower edges evenly rounded and provided with setae. Side-plate 4 deepest, with upper half of posterior border rather deeply emarginate, and lower border and lower half of posterior border provided with setae.




Figure 4. *Synurella johanseni*, n. sp.

1 Head and antennae. 2 Accessory flagellum enlarged. 3 Maxilla 1. 4 Maxilla 2. 5 Maxillipeds. 6 Lower lip. 7 Upper lip. 8 Mandible. 9 Seta of molar enlarged.

Mandible; 3rd joint of palp a little shorter than 2nd, armed at apex with 4 or 5 long bristles and at the upper part of the front edge with a row of fine bristles; front edge of 2nd joint with 5 or 6 long bristles. Molar projecting considerably from surface of mandible, and provided on inner edge of grinding surface with a very finely pinnate seta which is as long as the molar; on edge opposite seta is a group of minute spines; grinding surface covered with fine, sharp teeth. Maxilla 1; inner plate with 6 or 7 plumose setae; outer plate with 7 stout spines, some of which are notched; palp with row of spines at apex, below which are scattered setules. Maxilla 2; inner plate with seta on apex and on inner margin; outer plates with setae at apex. Maxillipeds; inner plates with notched spines at apex; outer plates with spines at apex and on inner edges. Lower lip normal.





Figure 5. *Synurella johanseni*, n. sp.

1 Gnathopod 1, left, inside view. 2 Notched spine of palm. 3 Gnathopod 2, left, inside view. 4 Side-plate 4

Figure 6. *Synurella johanseni*, n. sp.

1 Peraeopod 5. 2 2nd and 3rd abdominal segments. 3 3rd abdominal segment showing slight production of lower posterior corner. 4 Uropod 1. 5 Uropod 2. 6 Telson and 3rd uropods. 7 3rd uropod.

Gnathopods short and strong. Gnathopod 1; joints 3-6 about as broad as long; 6th joint with sides parallel; palm nearly transverse, slightly convex and provided with a double row of notched spines and a few bristles. Dactyl reaching to end of palm. Gnathopod 2; 6th joint longer than broad and slightly widening distally; palm oblique, evenly convex and provided with a double row of notched spines and a few bristles; dactyl reaching to end of palm.

Pereopods slender; 1st and 2nd shorter than rest; 4th longest, proportionately longer in ♂; 3rd, 4th, and 5th pereopods with 2nd joint moderately expanded and bearing shallow serrations on posterior border; dactyls each bearing a setule on the inner edge near extremity.

Gnathopod 2, and pereopods 1-3 each provided with a single lamellar branchia; pereopods 4 and 5 each provided with two cylindrical branchiæ; pleon segment 1 in ♀ with a single, small, cylindrical branchia on each side.

Uropod 1 longest; peduncle a little longer than the subequal rami. Uropod 2 with peduncle as long as the subequal rami. Uropod 3 and telson in their normal position project at right angles to the urosome; uropod 3 in this position not reaching end of telson; peduncle broad and flat, and the single ramus small and triangular with 2 stout spines on outer margin; no spines on peduncle.

Telson as broad as long; the slightly convex sides somewhat converging; end emarginate, depression reaching about one-third length of telson; lobes each provided at apex with 4 or 5 stout spines.

Length.—6 mm.

Remarks.—To the middle of the ventral surface of each of the 2nd, 3rd and 4th thoracic segments is attached an elongated, papilliform process about half the length of the branchiæ. Appendages probably of a similar nature were first observed and described by G. O. Sars in 1867 in the fresh water species *Gammaracanthus lacustris* Sars and *Pontoporeia affinis* Lindström. S. I. Smith in 1874 also observed them in a species of *Pontoporeia* from the Great Lakes. These appendages, the function of which is not known, have apparently been observed only in fresh water species.

This is the first appearance of the genus *Synurella* in America; the two other species of the genus having been found in Germany and Russia. The closely related genus *Borula* was discovered in Hungary so that in all probability these genera will be found to occur throughout Russia, Siberia and northern North America.

43. *Weyprechtia pinguis* (Krøyer).

1838. *Gammarus pinguis* KRØYER, Danske Selsk. Afh., vol. 7, p. 252, pl. I, f. 5.

1906. *Weyprechtia pinguis* STEBBING, Tierreich, Amph. I, p. 382, and synonymy.

1909. *Weyprechtia pinguis* BRÜGGEN, Mém. Acad. Imp. Sci. St. Pétersbourg, sér. 8, vol. 18, No. 16, p. 36.

Station 41c: Bernard harbour, Northwest Territories (outer harbour), July 28, 1915, 5 fathoms, sandy mud with many algæ; 1 specimen.

Station 41u: Bernard harbour, Northwest Territories, end of August, 1915, from stomach of *Salvelinus malma* Walb.; 16 specimens.

Colour.—Dorsal parts of head and body segments dark vinaceous-brown. Sides of body segments and side-plates mottled with dark vinaceous brown and light grayish vinaceous. Antennæ striped with bordeaux and light grayish vinaceous. Gnathopods, pereopods, pleopods, uropods, and telson light grayish vinaceous.

Distribution.—Arctic ocean (circumpolar).

Family PHOTIDÆ.

44. *Protomedea fasciata* Krøyer.1842. *Protomedea fasciata* KRØYER, Naturh. Tidsskr., vol. 4, p. 154.1906. *Protomedea fasciata* STEBBING, Tierreich, Amph. I, p. 623, and synonymy.1913. *Protomedea fasciata* STEPHENSEN, Meddel. om Grønland, vol. XXII, p. 206.

Station 43b: Dolphin and Union strait (off Staphylton bay), Northwest Territories, 25-30 fathoms, sandy mud with pebbles, but no algæ; 1 specimen.

Distribution.—Arctic ocean, North Atlantic, North sea, Skagerrak, Kattegat, Greenland, Spitzbergen, Iceland, Finmark (Norway), Sweden, Denmark.

Family AMPITHOIDÆ.

45. *Ampithoe rubricata* (Montagu).1808. *Cancer* (*Gammarus*) *rubricatus* MONTAGU, Tr. Linn. Soc. London, vol. 9, p. 99, pl. 5, f. 1.1906. *Ampithoe rubricata* STEBBING, Tierreich, Amph. I, p. 639, and synonymy.

Station 14: Lat. 54° 23' N., long. 164° 45' W., July 2, 1913, surface; 1 specimen.

Station 20b, c: Grantley harbour (port Clarence), Alaska, July 30, 1913, 2-3 fathoms, sandy mud with algæ; 1 specimen.

Distribution.—North Atlantic, with adjoining seas (Europe).

These records greatly extend the range of this species northward and westward.

An immature specimen of some species of *Ampithoe* was obtained at Station 20a, Grantley harbour, Alaska.

Family ISCHYROCERIDÆ.*

46. *Ischyrocerus anguipes* Krøyer.1838. *Ischyrocerus anguipes* KRØYER, Danske Selsk. Afh., vol. 7, p. 283, pl. 3, f. 14 a-m.1906. *Ischyrocerus anguipes* STEBBING, Tierreich, Amph. I, p. 658, and synonymy.1913. *Ischyrocerus anguipes* TATTERSALL, Proc. Roy. Irish Acad., vol. XXXI, Part 42, p. 18.1916. *Ischyrocerus anguipes* STEPHENSEN, Meddel. om Grønland, vol. LIII, p. 294.

Station 37j: Bernard harbour, Northwest Territories, September 1, 1914, pelagic, over 2 fathoms of water; 6 specimens.

Station 41s: Bernard harbour, Northwest Territories (inner harbour), August 24, 1915, surface; 1 specimen.

Station 41u: Bernard harbour, Northwest Territories, end of August, 1915, from stomach of *Salvelinus malma* Walb.; 1 specimen.

Station 42z: Bernard harbour, Northwest Territories (Dolphin and Union strait), December 12, 1915 (midnight), 0-3 fathoms; 1 specimen.

Distribution.—Arctic ocean (widely distributed), North Atlantic, North sea, Norway, West Baltic.

Family COROPHIIDÆ.

47. *Corophium bonellii* M.-Edw.1830. *Corophia bonellii* H. MILNE EDWARDS, Ann. Sci. Nat., vol. 20, p. 385.1906. *Corophium bonellii* STEBBING, Tierreich, Amph. I, p. 691, and synonymy.

Station 20b, c: Grantley harbour (port Clarence), Alaska, July 30, 1913, 2-3 fathoms, sandy mud with many algæ; 5 specimens.

Five specimens (1 male and 4 females) which appear to be *Corophium bonellii* were taken at Grantley harbour, Alaska. The females agree quite

* I have used the family name Ischyroceridæ rather than Jansidæ as the latter was created by Fieber in 1866 for a family of Hemiptera.

closely with Sars's figure. As the male has not been described, I give here a description of the specimen taken on this expedition.

Head with rostrum long and spear-shaped; lateral lobes short, apically rounded. Eyes dark, not very prominent.

Antenna 1 reaching about middle of 5th joint of antenna 2; 1st joint of peduncle not very thick, flattened dorsally and armed below with one terminal spine and one near centre; 2nd joint thinner and a little shorter, 3rd joint a little thinner than 2nd and not quite half as long; flagellum about $\frac{3}{4}$ length of peduncle and composed of 7 joints. Antenna 2; 2nd joint reaching very little beyond lateral lobes of head; 3rd joint equal in length to 2nd; 4th joint thickened and equal in length to 5th joint plus flagellum; lower distal corner of 4th joint produced into a strong, forward-pointing tooth above which is a much shorter one; 5th joint slender and slightly curved, inner distal edge produced into a blunt rounded lobe; no tooth on under side of 5th joint; flagellum composed of 3 joints, the last very short, bearing 2 curved spines and several setae.

Gnathopods, pereopods, uropods, and telson as in female.

Length.—4 mm.

Distribution.—North Atlantic, North sea, Skagerrak, and English channel, Norway, West France.

The present record marks a considerable western extension of the range of this species.

Family PODOCERIDÆ.

48. *Dulichia porrecta* (Bate).

1857. *Dyopodos porrectus* BATE, Ann. Nat. Hist., ser. 2, vol. 19, p. 151.

1906. *Dulichia porrecta* STEBBING, Tierreich, Amph. I, p. 712, and synonymy.

1913. *Dulichia porrecta* STEPHENSEN, Meddel. om Grønland, vol. XXII, p. 218.

Station 57a: Cape Smyth (point Barrow), Alaska, August 8, 1916, pelagic, over 1 fathom water; 1 specimen.

Distribution.—Arctic ocean, North Atlantic, North sea, West Greenland, Iceland, Lofoten island, South and West Norway, Danish waters, Shetlands.

SUB-ORDER CAPRELLIDEA.

Family CAPRELLIDÆ.

49. *Caprella drepanochir* Mayer.

1890. *Caprella drepanochir* MAYER, F. Fl. Neapel, vol. XVII, p. 81, pl. 7, f. 15, 33-34.

1903. *Caprella drepanochir* MAYER, Siboga-Exped., vol. XXXIV, p. 100, pl. 4, f. 11.

Station 20g: Port Clarence, Alaska, August 4, 1913, 2-3 fathoms, mud with many algæ; 7 specimens.

Station 20h: Port Clarence, Alaska, August 4, 1913, surface (attached to floating algæ); 20 specimens.

Distribution.—Collected between China and Mouth of Amur river, Vladivostok, Bering island, Chamisso harbour, and Eschscholtz bay, Alaska.

SUB-ORDER HYPERIIDÆ.

Family HYPERIIDÆ.

50. *Euthemisto libellula* (Mandt).

1822. *Gammarus libellula* MANDT, Obs. Hist. nat. et Anat. comp. in itinere groenlandico factae, p. 32.

1895. *Euthemisto libellula* SARS, Crust. Norway, vol. I, p. 13, pl. 6, f. 1.

Station 29f: Lat. 70° 13' N., long. 140° 50' W., April 4, 1914, water depth about 30 fathoms, from stomach of *Phoca hispida* Schreber; 8 specimens.

Station 41u: Bernard harbour, Northwest Territories, end of August, 1915, from stomach of *Salvelinus malma* Walb.; 30 specimens.

Station 42h: Bay at Bernard harbour, Northwest Territories, September 22, 1915, beachwater; 5 specimens.

Station 43b: Dolphin and Union strait (off Stapyhton bay), Northwest Territories, September 14, 1915, 25-30 fathoms, sandy mud with pebbles, but no alga; 7 specimens.

Station 66a: Latitude about 73° 50' N., long. 150° 15' W., August 31, 1918: surface; S. Storkerson, collector; 3 specimens.

Station 66b: Latitude about 73° 50' N., long. 147° W., September 17, 1918: surface; S. Storkerson, collector; 6 specimens.

Distribution: Arctic ocean, Greenland, Spitzbergen, Nova Scotia, Norway, Nova Zembla.

51. *Hyperia galba* (Montagu).

1813. *Cancer gammarus galba* MONTAGU, Trans. Linn. Soc., vol. XI, p. 4, pl. 2, f. 2.

1895. *Hyperia galba* Sars, Crust. Norway, vol. I, p. 7, pl. 2; pl. 3, f. 1, and synonymy.

Station 9a: Lat. 55° 2' N., long. 144° W., June 27, 1913, surface; 2 specimens.

Station 27h: Lagoon-bay at Collinson point, Alaska, September 18, 1913, 0-1 foot of water; 1 specimen.

Station 27m: Collinson point, Alaska, September 19, 1913, pelagic, over 1 foot of water (9-inch ice); 1 specimen.

Station 27u: Collinson point, Alaska, October 5, 1913, pelagic, over 1 fathom of water; 1 specimen.

Station 27y: Lagoon at Collinson point, Alaska, October 8, 1913, pelagic, over 2 feet of water; 1 specimen.

Station 30a: Lat. 69° 41' N., long. 141° 11' W., May 4, 1915, pelagic, over 3 fathoms of water; 1 specimen.

Color.—Animal translucent with dark vinaceous-drab markings, eyes very large and black, thorax with broad, dark vinaceous-drab band on side, second and third abdominal segments with dark vinaceous-drab dorsal spots, distal ends of the second joints of the gnathopods and first, second and third peraeopods marked with vinaceous-drab, peduncles of the pleopods also marked with vinaceous-drab.

Distribution.—Atlantic coast of France and Britain, Baltic, Arctic ocean, Greenland, Spitzbergen, Nova Zembla, Kara sea, Murman coast.

52. *Hyperoche kroeyeri* Bovallius.

1885. *Hyperia kroeyeri* BOVALLIUS, K. Svenska Vet.-Akad. Handlingar, Band 10, No. 14, p. 17.

1887. *Hyperoche kroeyeri* BOVALLIUS, *ibid.*, Band XI, No. 16, p. 18.

1895. *Hyperoche kroeyeri* Sars, Crust. Norway, vol. I, p. 9, pl. 4, and synonymy.

Station 27h: Lagoon-bay at Collinson point, Alaska, September 18, 1913, 0-1 foot of water; 1 specimen.

Station 27m: Collinson point, Alaska, September 19, 1913, pelagic, over 1 foot of water (9-inch ice); 12 specimens.

Station 27n: Collinson point, Alaska, September 20, 1913, pelagic, over 1 foot of water (9-inch ice); 6 specimens.

Station 57a: Cape Smyth (Point Barrow), Alaska, August 8, 1916, pelagic, over 1 fathom of water; 4 specimens.

Colour.—Central areas of eyes duck green. Dorsal parts of the body segments apricot orange. Sides of body, sideplates, gnathopods, peraeopods, pleopods, uropods and telson splotched with apricot orange. Rest of animal translucent.

Mr. Johansen states that this species was found symbiotic in a large etenophore.

Distribution.—Arctic ocean, Greenland, Labrador, Spitzbergen, White sea, Siberian polar sea.

53. *Parathemisto oblivia* (Krøyer).

1836. *Hyperia oblivia* KRØYER, Danske Vid. Selsk. Afhandl. vol. 7, p. 70, pl. 4, f. 19.

1895. *Parathemisto oblivia* Sars, Crust. Norway, vol. I, p. 10, pl. 5, f. 1, and synonymy.

Station 13g, h: Lat. $54^{\circ} 30' N.$ long. $159^{\circ} 42' W.$, July 1, 1913, surface; 1 specimen.

Station 14: Lat. $54^{\circ} 23' N.$, long. $164^{\circ} 45' W.$, July 2, 1913, surface; 2 specimens.

Station 21c: Lat. $68^{\circ} 48' N.$, long. $165^{\circ} 10' W.$, August 16, 1913, surface; 2 specimens.

Station 27g: Collinson point, Alaska, September 26, 1913, pelagic, over 1 fathom of water; 1 specimen.

Distribution.—British Isles, Greenland, Norway, Nova Zembla, Barents sea, North Atlantic, East coast of United States.

About 50 immature Hyperiidids were collected at station 18a, c, e, lat. $62^{\circ} N.$, long. $167^{\circ} 30' W.$, at the surface.

APPENDIX.

Additional data for the report upon the Amphipods of the Canadian Arctic Expedition, based upon specimens from the *Neptune* and other Canadian Expeditions.

BY CLARENCE R. SHOEMAKER.

Family LYSIANASSIDÆ.

1. *Anonyx nugax* (Phipps)

Cumberland gulf, Northwest Territories, September 4, 1904, from stomach of *Cottus (Myoxocephalus) groenlandicus* Bean, *Neptune* expedition; 6 specimens.
Cape Fullerton, west side Hudson bay, *Neptune* expedition, 1903-4; 6 specimens.

Hudson bay or strait, 1897? *Diana* expedition; 7 specimens.

Winter harbour, Melville island, Northwest Territories, *Arctic* expedition, 1909; 1 specimen.

Near mouth of Povungnituk river, east side of Hudson bay, Northwest Territories, 1898; A. P. Low, collector; 2 specimens.

2. *Onisimus edwardsii* (Krøyer).

1846. *Anonyx edwardsii* KRØYER, Naturh. Tidsskr., ser. 2, vo 12, p. 1, 41.

1906. *Onisimus edwardsii* STEBBING, Das Tierreich, Amph. I, p. 25, and synonymy.

1912. *Onisimus edwardsii* STEPHENSEN, Meddel. om Grønland, vol. XLV, p. 530.

1913. *Onisimus edwardsii* STEPHENSEN, Meddel. om Grønland, vol. XXII, p. 121.

1916. *Onisimus edwardsii* STEPHENSEN, Meddel. om Grønland, vol. LIII, 1916, p. 285.

Cape Fullerton, west side of Hudson bay, *Neptune* expedition, 1903-4; 5 specimens.

Distribution.—Arctic ocean, North Atlantic and North sea, West Norway.

3. *Orchomenella pinguis* (Boeck).

1861. *Anonyx pinguis* BOECK, Forh. Skand. Naturf., møde 8, p. 642.

1906. *Orchomenella pinguis* STEBBING, Das Tierreich, Amph. I, p. 82, and synonymy.

1913. *Orchomenella pinguis* STEPHENSEN, Meddel. om Grønland, vol. LI, p. 66.

1916. *Orchomenella pinguis* STEPHENSEN, Meddel. om Grønland, vol. LIII, p. 286.

Cape Fullerton, west side Hudson bay, *Neptune* expedition, 1903-4; 3 specimens.

Distribution.—Arctic ocean, North Atlantic, North sea and Skagerrak, Siberia, South and West Norway, Malangen fjord, Finland, Mediterranean.

4. *Socarnes bidenticulatus* (Bate).

Winter harbour, Melville island, Northwest Territories, *Arctic* Expedition. May 13, 1909; 1 specimen.

Winter harbour, Melville island, Northwest Territories, 7 fathoms. May 28, 1909, *Arctic* Expedition; 2 specimens.

Family AMPELISCIDÆ.

5. *Ampellicca eschrichtii* Krøyer.

Port Burwell, Ungava, July 28, 1904, *Neptune* expedition, 1903-4; 6 specimens.

6. *Byblis* species.

Port Burwell, Ungava, July 28, 1904, *Neptune* expedition, 1903-4; 1 specimen.

The poor condition of this specimen does not warrant a specific identification.

Family ACANTHONOTOZOMATIDÆ.

7. *Acanthonotosoma serratum* (Fabricius).

1780. *Oniscus serratus* FABRICIUS, Fauna Groenl., p. 262.

1906. *Acanthonotosoma serratum* STEBBING, Das Tierreich, Amph. I, p. 218.

1912. *Acanthonotosoma serratum* STEPHENSEN, Meddel. fra den Naturh. Foren., vol. 64, p. 93.

1912. *Acanthonotosoma serratum* STEPHENSEN, Meddel. om Grønland, vol. XIV, p. 596.

1913. *Acanthonotosoma serratum* STEPHENSEN, Meddel. om Grønland, vol. XXII, p. 167.

1913. *Acanthonotosoma serratum* STEPHENSEN, Meddel. om Grønland, vol. LI, p. 67.

1916. *Acanthonotosoma serratum* STEPHENSEN, Meddel. om Grønland, vol. LIII, p. 289.

Port Burwell, Ungava, July 28, 1904, *Neptune* expedition; 2 specimens.

Distribution.—Arctic ocean, North Atlantic, North sea and Skagerrak, Greenland, Spitzbergen, Barents sea, Murman coast, Kara sea, North America, Bohuslän, Norway from Haugesund northward.

Family OEDICEROTIDÆ.

8. *Paroediceros lynceus* (M. Sars).

Port Burwell, Ungava, July 28, 1904, *Neptune* expedition; 1 specimen.

Family PONTOGENEIIDÆ.

9. *Pontogeneia inermis* (Krøyer).

Cape Fullerton, west side Hudson bay, *Neptune* expedition, 1903-4; 1 specimen.

Family ATYLIDÆ.

10. *Atylus carinatus* (Fabricius).

Winter harbour, Melville island, Northwest Territories, *Arctic* expedition, 1907-9; 3 specimens.

Winter harbour, Melville island, Northwest Territories, *Arctic* expedition, May, 1909; 1 specimen.

Winter harbour, Melville island, Northwest Territories, *Arctic* expedition, 7 fathoms, May 28, 1909; 1 specimen.

Family EUSIRIDÆ.

11. *Rhachotropis aculeata* (Lepechin).

Hudson strait? Labrador? Hudson bay? *Neptune* expedition, 1903-4; 2 specimens.

Family GAMMARIDÆ.

12. *Gammarus locusta* (Linné).

Winter harbour, Melville island, Arctic expedition, May, 1909?; 8 specimens.

Winter harbour, Melville island, Northwest Territories, Arctic expedition, 7 fathoms, May 28, 1909; 1 specimen.

Wakeham bay, south side of Hudson strait, Ungava, September, 1904, Neptune expedition; 13 specimens.

13. *Gammaracanthus loricatus* (Sabine).

Winter harbour, Melville island, Northwest Territories, Arctic expedition; 1 specimen.

Winter harbour, Melville island, Northwest Territories, 7 fathoms, May 28, 1909, Arctic expedition; 1 specimen.

Winter harbour, Melville island, Northwest Territories, May, 1909?, Arctic expedition; 1 specimen.

Family ISCHYROCERIDÆ.

14. *Ischyrocerus anguipes* Kröyer.

Cape Fullerton, west side of Hudson bay, Northwest Territories, Neptune expedition, May 27, 1904 (Beach?); 1 specimen.

Family HYPERIIDÆ.

15. *Ethemisto compressa* (Goës).

1865. *Ethemisto compressa* Goës, Öfvers. af Kgl. Svenska Vetensk.-Akad. Forhandl., p. 533, pl. 41, fig. 34.

1870. *Parathemisto compressa* BOECK, Crust. Amph. boreal. et arct., Vid. Selsk. Forh. Christiania, p. 7.

1895. *Ethemisto compressa* G. O. Sars, Crustacea of Norway, vol. I, p. 12, pl. 5, f. 2.

1912. *Ethemisto compressa* STEPHENSEN, Meddel. fra den naturh. Foren., vol. 64, p. 84.

1912. *Ethemisto compressa* STEPHENSEN, Meddel. om Grønland, vol. XLV, p. 613.

1913. *Ethemisto compressa* STEPHENSEN, Meddel. om Grønland, vol. XXII, p. 103.

1913. *Ethemisto compressa* TATTERSALL, Proc. Royal Irish Academy, vol. XXXI, pt. 42, p. 21.

1916. *Ethemisto compressa* STEPHENSEN, Meddel. om Grønland, vol. LIII, p. 275.

Black Tickle, Labrador, beginning of September, 1903, pelagic, Neptune expedition; 1 specimen.

Distribution.—Arctic ocean, Davis strait, east coast of Greenland, New England coast, Norwegian coast.

16. *Ethemisto libellula* (Mandt).

Cumberland gulf, east of Blacklead island, (Baffin island), September 4, 1904, Neptune expedition, from stomach of *Cottus* (*Myxocephalus*) *groenlandicus* Bean; 2 specimens.

ADDITIONAL NOTE.

Some of the Amphipod Crustaceans collected by the Neptune Expedition, 1903-04, were sent to Prof. G. O. Sars, Christiania, Norway, for identification and are published (p. 368) in the report of the Department of Marine and Fisheries, Ottawa, Canada, 1905 (1906).

As these specimens are still in Christiania they have not been examined at the United States National Museum, Washington, nor included in the report above.

Professor Sars's determinations follow:—

- Anonyx nugax* (Phipps), Fullerton, Northwest Territories.
Pseudolibrotus littoralis (Krøyer), Fullerton, Northwest Territories.
Ischyrocerus angripes (Krøyer), juv., Fullerton, Northwest Territories.
Ampelisca eschrichti Krøyer, Port Burwell, Ungava.
Euthemisto libellula (Mandt), North Somerset, Northwest Territories.
Gammarus locusta (Linn.), Wakeham bay, Ungava.

BIBLIOGRAPHY.

- Birula, A.
 Researches in the Biology and Zoogeography of the Principal Seas of Russia. II. Hydrozoa, Polychaeta and Crustacea, collected by Dr. A. Botkine in 1895 from the Gulfs of the Yenisei and the Obi. Annuaire du Musée Zoologique de l'Académie Impériale des Sciences de St. Pétersbourg, vol. II, 1897, p. 78.
 Researches etc. VII. Note on the Crustacea collected by Dr. A. S. Botkine in 1896 and 1897 from the Kara Sea and the Region southeast of the Murman Sea. L. c., vol. IV, 1900, p. 418.
- Blanc, Henri.
 Die Amphipoden der Kieler Bucht nebst einer histologischen Darstellung der "Calceoli". Nova Acta der Kgl. Leop.-Carol. Deutschen Akademie der Naturforscher, vol. XLVII, 1884, Nr. 2.
- Bovallius, Carl.
 Some Forgotten genera of Amphipoda. K. Svenska Vet.-Akad. Handlingar, vol. 10, 1885, No. 14.
 Systematical List of the Amphipoda Hyperideae. L. c., vol. II, 1887, No. 16.
- Bradley, J. Chester.
 Notes on the Amphipods of the Genus *Corophium* from the Pacific Coast. University of California Publications in Zoology, vol. 4, No. 4, May 15, 1908.
- Brüggen, Ernest von der.
 Beiträge zur Kenntnis der Amphipoden-Fauna der russischen Arctis. Mémoires de l'Académie Impériale des Sciences de St. Pétersbourg, Sér. 8, Classe Physico-Mathématique, vol. XVIII, 1909, No. 16.
- Buchholz, R.
 Die zweite deutsche Nordpolarfahrt in den Jahren 1869 und 1870 unter Führung des Kapitan Karl Koldewey. Herausgegeben von dem Verein für die deutsche Nordpolarfahrt in Bremen. Vol. 2, Wissenschaftliche Ergebnisse. 8, Crustaceen. Leipzig. 1874.
- Chevreaux, Ed.
 Description d'un Amphipode (*Katius obesus*, nov. gen. et sp.), suivie d'une liste des Amphipodes de la tribu des Gammarina ramenés par le filet à grande ouverture pendant la dernière campagne de la *Princesse-Alice* en 1904. Bulletin du Musée Océanographique de Monaco, no. 35, May 5, 1905.
- Ekman, Sven.
 Zwei neue europäische Arten der Amphipodengattung *Pontoporeia* Krøyer. Arkiv för Zoologi utgivet af K. Svenska Vetenskapsakademien i Stockholm, vol. 8, 1913, No. 8.
- Goës, A.
 Crustacea amphipoda maris Spitsbergiam alluentis, cum speciebus salinis arcticis enumerat. Översigt Kongl. Vetenskaps-Akademien. Forhandlingar, Årg. 22, No. 8, Stockholm, 1865.
- Hansen, H. J.
 Malacostraca marina Groenlandiae occidentalis. Oversigt over det vestlige Grønlands Fauna af Malakostrake Havkrebsdyr. Vidensk. Meddel. fra den naturh. Foren. i Kjøbenhavn 1887.
 "Dijmphna"-Togtets zoologisk-botaniske Udbytte. Udgivet af Kjøbenhavns Universitets zoologiske Museum ved Chr. Fr. Lütken, pp. 157-286; pls. 18-24. Copenhagen, 1887.
- Holmes, S. J.
 The Amphipods collected by the U.S. Bureau of Fisheries Steamer *Albatross* off the West Coast of North America, in 1903 and 1904, with description of a new Family and several new Genera and Species. Proc. United States National Museum, vol. XXXV, 1908, pp. 489-543.
 Amphipod Crustaceans of the Harriman Alaska Expedition. Harriman Alaska Expedition, vol. 10, 1904, pp. 233-246.
- Mayer, P.
 Fauna und Flora des Golfes von Neapel und der angrenzenden Meeres-Abschnitte, herausgegeben von der zoologischen Station zu Neapel, vol. VI, 1882. Caprelliden.
 Die Caprellidae der Siboga-Expedition. Siboga-Expedition, Monographie XXXIV, 1903.

Ohlin, Axel.

Bidrag till Kännedomen om Malakostrakfaunan i Baffin Bay och Smith Sound. Acta Universitatis Lundensis. Lunds Universitets Års-Skrift., vol. XXXI., 1895, pp. 20-65; plate.

Pearse, A. S.

Notes on a small collection of Amphipods from the Pribilof Islands, with Descriptions of New Species. Proc. United States National Museum, vol. 45, 1913, pp. 571-573.

Sars, G. O.

Crustacea I, II. The Norwegian North Atlantic Expedition, 1876-1878. Zoology, vols. XIV and XV, 1885.

An account of the Crustacea of Norway, with short Descriptions and Figures of all the Species. vol. I, 1895, text and plates.

The Norwegian North Polar Expedition 1893-1896. vol. I, 1900, Part V. Crustacea.

Report of the Second Norwegian Arctic Expedition in the "Fram" 1898-1902, vol. III, 1909, Crustacea, No. 18, pp. 14-34, pls. 1-6.

Stappers, Louis.

Crustacés Malacostracés. Campagne Arctique de 1907, du Due d'Orléans. 1911.

Stebbing, T. R. R.

Das Tierreich. Amphipoda. I, Gammaridea. Berlin, 1906.

Stephensen, K.

Report on the Malacostraca Pycnogonida and some Entomostraca collected by the "Danmark" Expedition to Northeast Greenland. Meddelelser om Grønland, vol. XLV, 1912, pp. 526-544.

Report on the Malacostraca collected by the "Tjalfe" Expedition, under the direction of cand. mag. Ad. S. Jensen, especially at W. Greenland. Meddel. fra den Naturh. Foren. i København, vol. 64, 1912, pp. 81-97.

Grønlands Krebsdyr og Pycnogonider (Conspectus Crustaceorum et Pycnogonidorum Groenlandiae). Meddelelser om Grønland, vol. XXII, 1913, pp. 93-228.

Account of the Crustacea and the Pycnogonida collected by Dr. V. Nordmann in the summer of 1911 from Northern Strömfjord and Giesecke Lake in West Greenland. Meddelelser om Grønland, vol. LI, 1913, pp. 65-68.

Zoogeographical Investigation of certain Fjords in Southern Greenland. Meddelelser om Grønland, vol. LIII, 1916, pp. 275-297.

Stuxberg, Anton.

In A. E. Nordenskiöld, Vega Expeditionens Vetenskapliga Iakttagelser, vol. I, 1882, pp. 679-812; vol. 5, 1887, pp. 60-73.

In A. E. Nordenskiöld, The Voyage of the Vega, vol. I, II, 1881.

Tattersall, W. M.

The Marine Fauna of the Coast of Ireland, Part VIII. Pelagic Amphipoda of the Irish Atlantic Slope. "Fisheries, Ireland, Sci. Invest., No. IV, 1905 [1906]", pp. 1-39, pls. 1-5.

Clare Island Survey, Part 42, Amphipoda. Proc. Royal Irish Academy, vol. XXXI, 1913, pp. 1-24.

Weckel, Ada L.

The Fresh-water Amphipods of North America. Proc. of the United States National Museum, vol. XXXII, pp. 25-58.

l. Acta
905, pp.

otions of
73.

y, vols.

all the

ol. III,

"Dan-
XLV,

tion of
Foren.

idorum

ummer
lelelser

ser om

2, pp.

e Irish
9, pls.

1913,

seum,

8-1-18